Author Index

ABDEL-MOTAL U.M. *see* FAYOLLE C.
ABKEN H. *see* REINHOLD U.
ABO T. *see* KAWAMURA T.
ABRAHAMSOHN I.A. *see* TADOKORO C.E.
ÁGUAS A.P. *see* BRÁS A.
AKBAR A.N. *see* GOMBERT W.
ALAVA M.A. *see* MARTÍNEZ-LORENZO M.J.
ALBERS R., VAN DER PIJL A., SEINEN W., PIETERS R. & BLOKSMA N. The autoimmunogenic chemicals $HgCl_2$ and diphenylhydantoin stimulate IgG production to TNP–Ficoll and TNP–OVA, supporting and extending the graft-versus-host hypothesis for chemical induction of autoimmunity, 468
ALCOCER J.M. *see* HERNÁNDEZ-PANDO R.
ALLEN R.F. *see* BARKER R.N.
ALTSTAEDT J., KIRCHNER H. & RINK L. Cytokine production of neutrophils is limited to interleukin-8, 563
ANEL A. *see* MARTÍNEZ-LORENZO M.J.
AOKI Y. *see* HIROMATSU K.
AOYAGI T. *see* MIGITA K.
APPELBERG R. *see* LEAL I.S.
ARAMAKI Y. *see* MURAI M.
ASNAGHI V. *see* MATERA L.
ATKINS C. *see* SANCHEZ-GARCIA J.
AVANZI G. *see* MATERA L.

BACCANTE G. *see* PORRECA E.
BAL V. *see* MENDIRATTA S.K.
BALL S. *see* FIDLER S.J.
BANDO Y. & HIGGINS D.A. Duck lymphoid organs: their contribution to the ontogeny of IgM and IgY, 8
BARBACANE R.C. *see* PORRECA E.
BARJA-FIDALGO C. *see* FIERRO I.M.
BARKER R.N. *see* SHEN C.-R.
BARKER R.N., EASTERFIELD A.J., ALLEN R.F., WELLS A.D., ELSON C.J. & THOMPSON S.J. B- and T-cell autoantigens in pristane-induced arthritis, 189
BARNES P.J. *see* BERKMAN N.
BEDFORD P.A. *see* COATES J.P.
BEECH J.T. *see* SHEN C.-R.
BEFUS A.D. *see* LIN T.J.
BELLONE G. *see* MATERA L.
BENEDETTI C. *see* ROSATI E.
BERG L. *see* FAYOLLE C.
BERGER P. *see* KLONISCH T.
BERKMAN N., ROBICHAUD A., ROBBINS R.A., ROESEMS G., HADDAD E.B., BARNES P.J. & CHUNG K.F. Inhibition of inducible nitric oxide synthase expression by interleukin-4 and interleukin-13 in human lung epithelial cells, 363
BEVERLEY P.C.L. *see* GOMBERT W.
BISSONNETTE E.Y. *see* LIN T.J.
BLOKSMA N. *see* ALBERS R.
BOFILL M. *see* GOMBERT W.
BONDER C.S. *see* HART P.H.
BONIFAS U. *see* FROSCH S.
BONIVER J. *see* DEMAN J.
BORGHESI C. & NICOLETTI C. Autologous anti-idiotypic antibody response is regulated by the level of circulating complementary idiotype, 172
BORREBAECK A.K. *see* DUEÑAS M.
BORTHWICK N.J. *see* GOMBERT W.
BOWLES V.M. *see* EGAN P.J.
BRANDON M.R. *see* EGAN P.J.
BRANDT E.R., HAYMAN W.A., CURRIE B., CARAPETIS J., WOOD Y., JACKSON D.C., COOPER J., MELROSE W.D., SAUL A.J. & GOOD M.F. Opsonic human antibodies from an endemic population specific for a conserved epitope on the M protein of group A streptococci, 331
BRANDTZAEG P. *see* FARSTAD I.N. *see also* HELGELAND L.
BRÁS A. & ÁGUAS A.P. Diabetes-prone NOD mice are resistant to *Mycobacterium avium* and the infection prevents autoimmune disease, 20
BRENNER B. *see* GULBINS E.
BROCHIER J. *see* GAILLARD J.P.
BRUSCO A. *see* ENGSTRÖM P.-E.

CALDERÓN G.M. *see* PÉREZ C.
CARAPETIS J. *see* BRANDT E.R.
CARBONARA A.O. *see* ENGSTRÖM P.-E.
CARRIÓ R. *see* RAMOS A.
CASALVILLA R. *see* DUEÑAS M.
CASTELLANI M.L. *see* PORRECA E.
CEUPPENS, J.L. *see* KRUGER M.
CHAN J. *see* LU J.
CHEONG C.-H. *see* JEONG S.-Y.
CHILVERS E.R. *see* CONDLIFFE A.M.
CHIN, L.-T. *see* DUEÑAS M.
CHULUYÁN E. *see* MORELLI A.
CHUNG K.F. *see* BERKMAN N.
CIANETTI S. *see* ROSATI E.
CLAASSEN E. *see* DEMAN J.
COATES J.P., ROWLAND S., HILL S., IQBALL S., BEDFORD P.A., KIMBER I. & KNIGHT S.C. Comparison between the phenotype and function of maturing dendritic cells from spleen and lymph nodes, 457
COCHRANE L. *see* HARPER H.M.
COGGESHALL K.M. *see* GULBINS E.
COLOMB M.G. *see* VILLIERS M.-B.
COLSTON M.J. *see* SANCHEZ-GARCIA J.
CONDLIFFE A.M., CHILVERS E.R., HASLETT C. & DRANSFIELD I. Priming differentially regulates neutrophil adhesion molecule expression/function, 105
CONTARINI M. *see* MATERA L.
CONTI P. *see* PORRECA E.
COOPER J. *see* BRANDT E.R.
COOREVITS, L. *see* KRUGER M.
CORATO A. *see* SHEN C.-R.
CORNACCHIONE P. *see* ROSATI E.
CUCCURULLO F. *see* PORRECA E.
CUNHA F.Q. *see* FIERRO I.M.
CURRIE B. *see* BRANDT E.R.

D'ORAZIO *see* LI X.-Y.
DE HAAN A., GROEN G., PROP J., VAN ROOIJEN N. & WILSCHUT J. Mucosal immunoadjuvant activity of liposomes: role of alveolar macrophages, 488
DE WITT, T.P.M. *see* KRUGER M.
DEARMAN R.J., MOUSSAVI A., KEMENY D.M. & KIMBER I. Contribution of $CD4^+$ and $CD8^+$ lymphocyte subsets to the cytokine secretion patterns induced in mice during sensitization to contact and respiratory chemical allergens, 502
DECK R.R. *see* ULMER J.B.
DEFRESNE M.P. *see* DEMAN J.
DELVES P.J. *see* KLONISCH T.
DEMAN J., VAN MEURS M., CLAASSEN E., HUMBLET C., BONIVER J. & DEFRESNE M.P. *In vivo* expression of interleukin-1β (IL-1β), IL-2, IL-4, IL-6, tumour necrosis factor-α and interferon-γ in the fetal murine thymus, 152
DERIAUD E. *see* FAYOLLE C.
DEVOS S. *see* MURAILLE E.

DEWITT C.M. *see* ULMER J.B.
DI FEBBO C. *see* PORRECA E.
DI GIOACCHINO M. *see* PORRECA E.
DÖBBELING U. The influence of IL-7 on V(D)J recombination, 569
DONNELLY J.J. *see* ULMER J.B.
DÖRMER P. *see* KHALIL R.M.A.
DORRELL L. *see* FIDLER S.J.
DRANSFIELD I. *see* CONDLIFFE A.M.
DRAPIER J.-C. *see* LE PAGE C.
DREXLER H.G. *see* QUENTMEIER H.
DUEÑAS M., CHIN, L.-T., MALMBORG A.-C., CASALVILLA R., OHLIN M. & BORREBAECK A.K. *In vitro* immunization of naive human B cells yields high affinity immunoglobulin G antibodies as illustrated by phage display, 1
DUSCHL A. *see* QUENTMEIER H.

EASTERFIELD A.J. *see* BARKER R.N.
EDWARDS P. *see* KLONISCH T.
EGAN P.J., KIMPTON W., SEOW H.-F., BOWLES V.M., BRANDON M.R. & NASH A.D. Inflammation-induced changes in the phenotype and cytokine profile of cells migrating through skin and afferent lymph, 539
EGUCHI K. *see* MIGITA K.
ELSON C.J. *see* BARKER R.N.
see also SHEN C.-R.
ENGSTRÖM P.-E., NORHAGEN E.G., OSIPOVA L., HELAL A., WIEBE V., BRUSCO A., CARBONARA A.O., LEFRANC G. & LEFRANC M.-P. Salivary IgG subclasses in individuals with and without homozygous IGHG gene deletions, 178
ENOSAWA S. *see* SUZUKI S.
ERNST M. *see* FLEISCHER J.

FAINBOIM L. *see* MORELLI A.
FARSTAD I.N., HALSTENSEN T.S., LIEN B., KILSHAW P.J., LAZAROVITZ A.I. & BRANDTZAEG P. Distribution of β_7 integrins in human intestinal mucosa and organized gut-associated lymphoid tissue, 227
FAYOLLE C., ABDEL-MOTAL U.M., BERG L., DERIAUD E., JONDAL M. & LECLERC C. Induction of cytotoxic T-cell response by optimal-length peptides does not require $CD4^+$ T-cell help, 41
FEARNLEY D.B. *see* WILLIAMS L.A.
FERGUSON A. Mucosal immunology: from bench to the bedside and beyond, 475
FERNANDEZ SUAREZ J.M. *see* GAILLARD J.P.
FERREIRA S.H. *see* FIERRO I.M.
FETTUCCIARI K. *see* ROSATI E.
FIDLER S.J., DORRELL L., BALL S., LOMBARDI G., WEBER J., HAWRYLOWICZ C. & REES A.D.M. An early antigen-presenting cell defect in HIV-1-infected patients correlates with CD4 dependency in human T-cell clones, 46
FIERRO I.M., BARJA-FIDALGO C., CUNHA F.Q. & FERREIRA S.H. The involvement of nitric oxide in the anti-*Candida albicans* activity of rat neutrophils, 295
FINLAY-JONES J.J. *see* HART P.H.
FLAD H.-D. *see* FLEISCHER J.
FLEISCHER J., SOETH E., REILING N., GRAGE-GRIEBENOW E., FLAD H.-D. & ERNST M. Differential expression and function of CD80 (B7–1) and CD86 (B7–2) on human peripheral blood monocytes, 592
FOA R. *see* MATERA L.
FRANCIA DI CELLE P. *see* MATERA L.
FRIEDMAN H. *see* RETZLAFF C.
FROSCH S., BONIFAS U., ROβ R., SCHWING J., YAGITA H., GUO Y., LIU Y. & RESKE-KUNZ A.B. Costimulatory signalling potential of murine MHC class II-positive T-clone cells, 384
FUJISAWA H., KONDO S., WANG B., SHIVJI G.M. & SAUDER D.N. The role of CD4 molecules in the induction phase of contact hypersensitivity cytokine profiles in the skin and lymph nodes, 250
FUKAMACHI H. *see* YANAGIDA M.

GABERT F.M. *see* VILLIERS M.-B.
GAILLARD J.P., LIAUTARD J., MANI J.C., FERNANDEZ SUAREZ J.M., KLEIN B. & BROCHIER J. Identification of a novel antigenic structure of the human receptor for interleukin-6 involved in the interaction with the glycoprotein 130 chain, 135
GAO J.-X. & ISSEKUTZ A.C. Expression of VCAM-1 and VLA-4 dependent T-lymphocyte adhesion to dermal fibroblasts stimulated with proinflammatory cytokines, 375
GASQUE P. & MORGAN B.P. Complement regulatory protein expression by a human oligodendrocyte cell line: cytokine regulation and comparison with astrocytes, 338
GOMBERT W., BORTHWICK N.J., WALLACE D.L., HYDE H., BOFILL M., PILLING D., BEVERLEY P.C.L., JANOSSY G., SALMON M. & AKBAR A.N. Fibroblasts prevent apoptosis of IL-2-deprived T cells without inducing proliferation: a selective effect on Bcl-x_L expression, 397
GONZÁLEZ M. *see* RAMOS A.
GOOD M.F. *see* BRANDT E.R.
GOODIER M.R. *see* MORRIS JONES S.
GRAGE-GRIEBENOW E. *see* FLEISCHER J.
GROEN G. *see* DE HAAN A.
GULBINS E., BRENNER B., SCHLOTTMANN K., WELSCH J., HEINLE H., KOPPENHOEFER U., LINDERKAMP O., COGGESHALL K.M. & LANG F. Fas-induced programmed cell death is mediated by a Ras-regulated O_2^- synthesis, 205
GUO Y. *see* FROSCH S.

HADDAD E.B. *see* BERKMAN N.
HAFIZI S. *see* MCDOUALL R.M.
HAGIWARA T. *see* YANAGIDA M.
HAIG D. McK., McINNES C.J., HUTCHISON G., SEOW H.-F. & REID H.W. Cyclosporin A abrogates the acquired immunity to cutaneous reinfection with the parapoxvirus orf virus, 524
HALSTENSEN T.S. *see* FARSTAD I.N.
HAMADA F. *see* HITSUMOTO Y.
HARADA M. *see* TAMADA K.
HARPER H.M., COCHRANE L. & WILLIAMS N.A. The role of small intestinal antigen-presenting cells in the induction of T-cell reactivity to soluble protein antigens: association between aberrant presentation in the lamina propria and oral tolerance, 449
HART D.N.J. *see* WILLIAMS L.A.
HART P.H., BONDER C.S., JONES C.A. & FINLAY-JONES J.J. Control of major histocompatibility complex class II expression on human monocytes by interleukin-4: regulatory effect of lipopolysaccharide, 599
HASHIMOTO K. *see* KAWAGUCHI-MIYASHITA M.
HASLETT C. *see* CONDLIFFE A.M.
HAWRYLOWICZ C. *see* FIDLER S.J.
HAYES K.C. *see* ISRAEL E.J.
HAYMAN W.A. *see* BRANDT E.R.
HEINLE H. *see* GULBINS E.
HELAL A. *see* ENGSTRÖM P.-E.
HELGELAND L., VAAGE J.T., ROLSTAD B., MIDTVEDT T. & BRANDTZAEG P. Microbial colonization influences composition and T-cell receptor Vβ repertoire of intraepithelial lymphocytes in rat intestine, 494
HERNÁNDEZ-PANDO R., OROZCOE H., SAMPIERI A., PAVÓN L., VELASQUILLO C., LARRIVA-SAHD J., ALCOCER J.M. & MADRID M.V. Correlation between the kinetics of Th1/Th2 cells and pathology in a murine model of experimental pulmonary tuberculosis, 26
HIGGINS D.A. *see* BANDO Y.
HILL S. *see* COATES J.P.
HINKULA J. *see* LUNDIN K.
HIROMATSU K., NISHIMURA H., KIMURA K., AOKI Y., USAMI J., KOBAYASHI N., MAKINO M. & YOSHIKAI Y. Th0-like $CD4^+$ T cells protect mice with murine retrovirus-induced immunodeficiency syndrome (MAIDS) against the co-infection with *Listeria monocytogenes*, 532
HIRSH A. *see* LIN T.J.
HITSUMOTO Y. *see* OKADA M.
HITSUMOTO Y., SONG D.-S., OKADA M., HAMADA F., SAHEKI S. & TAKEUCHI N. Enhancement of CD3-mediated thymocyte apoptosis by the cross-linkage of heat-stable antigen, 200

HODSON D. & OLIVEIRA D.B.G. The strain difference in the effect of mercuric chloride on antigen-triggered serotonin release from rat mast cells is not mediated via interferon-γ, 463
HOFFMAN P.S. *see* RETZLAFF C.
HONG S.-H. *see* JEONG S.-Y.
HOOGHE PETERS E.L. *see* MATERA L.
HORIBE K. *see* KIYOI H.
HORIUCHI T. *see* UEDA Y.
HU Z.-B. *see* QUENTMEIER H.
HÜLTNER L. *see* KHALIL R.M.A.
HUMBLET C. *see* DEMAN J.
HUTCHISON G. *see* HAIG D. McK.
HYDE H. *see* GOMBERT W.

ICHINOSE Y. *see* MIGITA K.
IIAI T. *see* KAWAMURA T.
IIKURA Y. *see* YANAGIDA M.
IKEDA K. *see* UEDA Y.
INOSHIMA Y. *see* PECORARO M.R.
IQBALL S. *see* COATES J.P.
ISHIKAWA H. *see* KAWAGUCHI-MIYASHITA M.
ISRAEL E.J., WILSKER D.F., HAYES K.C., SCHOENFELD D. & SIMISTER N.E. Increased clearance of IgG in mice that lack β_2-microglobulin: possible protective role of FcRn, 573
ISSEKUTZ A.C. *see* GAO J.-X.
ITO O. *see* TAMADA K.

JACKSON A.M. *see* KLONISCH T.
JACKSON D.C. *see* BRANDT E.R.
JACQUIER-SARLIN M.R. *see* VILLIERS M.-B.
JANOSSY G. *see* GOMBERT W.
JANSSON M. *see* LUNDIN K.
JEONG S.-Y., LEE J.-H., KIM H.-S., HONG S.-H., CHEONG C.-H. & KIM I.-K. 3-deazaadenosine analogues inhibit the production of tumour necrosis factor-α in RAW264.7 cells stimulated with lipopolysaccharide, 558
JONDAL M. *see* FAYOLLE C.
JONES C.A. *see* HART P.H.
JOURNET A.M. *see* VILLIERS M.-B.

KAI C. *see* PECORARO M.R.
KANEDA K. *see* ONODERA S.
KARDINAL C., SELMAYR M. & MOCIKAT R. Genetic stability of gene targeted immunoglobulin loci. I. Heavy chain isotype exchange induced by a universal gene replacement vector, 309
KATEVUO K. & VAINIO O. Thymocyte emigration in the chicken: an over-representation of $CD4^+$ cells over $CD8^+$ in the periphery, 419
KAWABE Y. *see* MIGITA K.
KAWACHI Y. *see* KAWAMURA T.
KAWAGUCHI Y. *see* PECORARO M.R.
KAWAGUCHI-MIYASHITA M., SHIMIZU K., NANNO M., SHIMADA S., WATANABE T., KOGA Y., MATSUOKA Y., ISHIKAWA H., HASHIMOTO K. & OHWAKI M. Development and cytolytic function of intestinal intraepithelial T lymphocytes in antigen-minimized mice, 268
KAWAMURA T., KAWACHI Y., MORODA T., WEERASINGHE A., IIAI T., SEKI S., TAZAWA Y., TAKADA G. & ABO T. Cytotoxic activity against tumour cells mediated by intermediate TCR cells in the liver and spleen, 68
KELLY P.A. *see* MATERA L.
KEMENY D.M. *see* DEARMAN R.J.
KHALIL R.M.A., HÜLTNER L., MAILHAMMER R., LUZ A., MOELLER J., MOHAMED A.A., OMRAN S. & DÖRMER P. Kinetics of interleukin-6 production after experimental infection of mice with *Schistosoma mansoni*, 256
KILSHAW P.J. *see* FARSTAD I.N.
see also PENNEY L.
KIM H.-S. *see* JEONG S.-Y.
KIM I.-K. *see* JEONG S.-Y.
KIMBER I. *see* COATES J.P.
see also DEARMAN R.J.
KIMPTON W. *see* EGAN P.J.
KIMURA K. *see* HIROMATSU K.
KIRCHNER H. *see* ALTSTAEDT J.
KIYOI H., MORI H., HORIBE K., OHNO R. & NAOE T. Comparison of the immunoglobulin heavy-chain complementarity determining region-3 structure among the DNA sequences and the μ- and γ-transcripts in human B-lineage cells, 324
KLEIN B. *see* GAILLARD J.P.
KLEIN T.W. *see* RETZLAFF C.
KLONISCH T., PANAYOTOU G., EDWARDS P., JACKSON A.M., BERGER P., DELVES P.J., LUND T. & ROITT I.M. Enhancement in antigen binding by a combination of synergy and antibody capture, 165
KNIGHT S.C. *see* COATES J.P.
KOBAYASHI N. *see* HIROMATSU K.
KODAMA T. *see* OKADA M.
KOGA Y. *see* KAWAGUCHI-MIYASHITA M.
KOLKOWSKI E. *see* MORELLI A.
KON O.L. *see* LU J.
KONDO S. *see* FUJISAWA H.
KOPPENHOEFER U. *see* GULBINS E.
KRUGER M., VAN DE WINKEL, J.G.J., DE WITT, T.P.M., COOREVITS, L. & CEUPPENS, J.L. Granulocyte–macrophage colony-stimulating factor down-regulates CD14 expression on monocytes, 89
KURIYAMA T. *see* ONODERA S.

LANG F. *see* GULBINS E.
LANGHORNE J. *see* MORRIS JONES S.
LARREGINA A. *see* MORELLI A.
LARRIVA-SAHD J. *see* HERNÁNDEZ-PANDO R.
LAZAROVITS A.I. *see* WALSH G.M.
see also FARSTAD I.N.
LE PAGE C., SANCEAU J., DRAPIER J.-C. & WIETZERBIN J. Differential expression of inducible NO synthase in two murine macrophage cell lines, 274
LE Y. *see* LU J.
LEAL I.S., APPELBERG R. & SILVA M.T. Neutrophils are involved in the non-specific resistance to listeriosis induced by mycobacterial infections, 442
LEBRUN J.-J. *see* MATERA L.
LECLERC C. *see* FAYOLLE C.
LEE J.-H. *see* JEONG S.-Y.
LEE S.H. *see* LU J.
LEFRANC G. *see* ENGSTRÖM P.-E.
LEFRANC M.-P. *see* ENGSTRÖM P.-E.
LEO O. *see* MURAILLE E.
LI X.-K. *see* SUZUKI S.
LI X.-Y., D'ORAZIO T. & NIEDERKORN J.Y. Role of Th1 and Th2 cells in anterior chamber-associated immune deviation, 34
LIAUTARD J. *see* GAILLARD J.P.
LIEN B. *see* FARSTAD I.N.
LIN T.J., BISSONNETTE E.Y., HIRSH A. & BEFUS A.D. Stem cell factor potentiates histamine secretion by multiple mechanisms, but does not affect tumour necrosis factor-α release from rat mast cells, 301
LINDERKAMP O. *see* GULBINS E.
LIU L. *see* REINHOLD U.
LIU M.A. *see* ULMER J.B.
LIU Y. *see* FROSCH S.
LOMBARDI G. *see* FIDLER S.J.
LÓPEZ-BOTET M. *see* PÉREZ-VILLAR J.J.
LÓPEZ-HOYOS M. *see* RAMOS A.
LU J., LE Y., KON O.L., CHAN J. & LEE S.H. Biosynthesis of human ficolin, an *Escherichia coli*-binding protein, by monocytes: comparison with the synthesis of two macrophage-specific proteins, C1q and the mannose receptor, 289
LUND T. *see* KLONISCH T.
LUNDIN K., SAMUELSSON A., JANSSON M., HINKULA J., WAHREN B., WIGZELL H. & PERSSON M.A.A. Peptides isolated from random peptide libraries on phage elicit a neutralizing anti-HIV-1 response: analysis of immunological mimicry, 579
LUZ A. *see* KHALIL R.M.A.

MACDONALD T.T. *see* PENNEY L.

MACEDO M.S. *see* TADOKORO C.E.
MADRID M.V. *see* HERNÁNDEZ-PANDO R.
MAEDA K. *see* PECORARO M.R.
MAILHAMMER R. *see* KHALIL R.M.A.
MAKINO M. *see* HIROMATSU K.
MALMBORG A.-C. *see* DUEÑAS M.
MANI J.C. *see* GAILLARD J.P.
MARCONI P. *see* ROSATI E.
MARIANO M. *see* MOURA A.C.N.
MARTÍNEZ-LORENZO M.J., ALAVA M.A., ANEL A., PIÑEIRO A. & NAVAL J. Release of preformed Fas ligand in soluble form is the major factor for activation-induced death of Jurkat T cells, 511
MATERA L., BELLONE G., LEBRUN J.-J., KELLY P.A., HOOGHE PETERS E.L., FRANCIA DI CELLE P., FOA R., CONTARINI M., AVANZI G. & ASNAGHI V. Role of prolactin in the *in vitro* development of interleukin-2-driven anti-tumoural lymphokine-activated killer cells, 619
MATSUNO T. *see* ONODERA S.
MATSUOKA Y. *see* KAWAGUCHI-MIYASHITA M.
MATSUZAKI G. *see* TAMADA K.
MAZZA G. *see* SHEN C.-R.
MCDOUALL R.M., PAGE C.S., HAFIZI S., YACOUB M.H. & ROSE M.L. Alloproliferation of purified $CD4^+$ T cells to adult human heart endothelial cells, and study of second-signal requirements, 220
McINNES C.J. *see* HAIG D. McK.
MCLELLAN A.D. *see* WILLIAMS L.A.
MELENDRO E.I. *see* PÉREZ C.
MELERO I. *see* PÉREZ-VILLAR J.J.
MELROSE W.D. *see* BRANDT E.R.
MENDIRATTA S.K., SINGH N., BAL V. & RATH S. Analysis of T-cell hybridomas with an unusual MHC class II-dependent ligand specificity, 238
MERINO J. *see* RAMOS A.
MEZZASOMA L. *see* ROSATI E.
MIDTVEDT T. *see* HELGELAND L.
MIGITA K., EGUCHI K., KAWABE Y., ICHINOSE Y., TSUKADA T., AOYAGI T., NAKAMURA H. & NAGATAKI S. TNF-α-mediated expression of membrane-type matrix metalloproteinase in rheumatoid synovial fibroblasts, 553
MIKAMI T. *see* PECORARO M.R.
MOCIKAT R. *see* KARDINAL C.
MOELLER J. *see* KHALIL R.M.A.
MOHAMED A.A. *see* KHALIL R.M.A.
MONROE J.G. *see* WECHSLER-REYA R.J.
MORELLI A., LARREGINA A., CHULUYÁN E., KOLKOWSKI E. & FAINBOIM L. Expression and modulation of C5a receptor (CD88) on skin dendritic cells. Chemotactic effect of C5a on skin migratory dendritic cells, 126
MORGAN B.P. *see* GASQUE P.
MORI H. *see* KIYOI H.
MORI T. *see* TAMADA K.
MORODA T. *see* KAWAMURA T.
MORRIS JONES S., GOODIER M.R. & LANGHORNE J. The response of γδ T cells to *Plasmodium falciparum* is dependent on activated $CD4^+$ T cells and the recognition of MHC class I molecules, 405
MOSER M. *see* MURAILLE E.
MOURA A.C.N. & MARIANO M. lipids from *Mycobacterium leprae* cell wall are endowed with an anti-inflammatory property and inhibit macrophage function *in vivo*, 613
MOUSSAVI A. *see* DEARMAN R.J.
MURAI M., ARAMAKI Y. & TSUCHIYA S. α_2-macroglobulin stimulation of protein tyrosine phosphorylation in macrophages via the mannose receptor for Fcγ receptor-mediated phagocytosis activation, 436
MURAILLE E., DEVOS S., THIELEMANS K., URBAIN J., MOSER M. & LEO O. Costimulation regulates the kinetics of interleukin-2 response to bacterial superantigens, 245
MURPHY F.J. & REEN D.J. Differential expression of function-related antigens on newborn and adult monocyte subpopulations, 587

NAGASAWA K. *see* UEDA Y.
NAGATAKI S. *see* MIGITA K.
NAKAHATA T. *see* YANAGIDA M.
NAKAMURA H. *see* MIGITA K.
NANNO M. *see* KAWAGUCHI-MIYASHITA M.
NAOE T. *see* KIYOI H.
NASH A.D. *see* EGAN P.J.
NAVAL J. *see* MARTÍNEZ-LORENZO M.J.
NEWTON S. *see* SHEN C.-R.
NICOLETTI C. *see* BORGHESI C.
NIEDERKORN J.Y. *see* LI X.-Y.
NIHO Y. *see* UEDA Y.
NISHIHIRA J. *see* ONODERA S.
NISHIMURA H. *see* HIROMATSU K.
NISHIZAKA H. *see* UEDA Y.
NOMOTO K. *see* TAMADA K.
NORHAGEN E.G. *see* ENGSTRÖM P.-E.

OHLIN M. *see* DUEÑAS M.
OHNO R. *see* KIYOI H.
OHWAKI M. *see* KAWAGUCHI-MIYASHITA M.
OKADA M. *see* HITSUMOTO Y.
OKADA M., SAGAWA T., TOMINAGA A., KODAMA T. & HITSUMOTO Y. Two mechanisms for platelet-mediated killing of tumour cells: one cyclo-oxygenase dependent and the other nitric oxide dependent, 158
OKUBO S. *see* RETZLAFF C.
OLIVEIRA D.B.G. *see* HODSON D.
OMRAN S. *see* KHALIL R.M.A.
ONODERA S., SUZUKI K., MATSUNO T., KANEDA K., KURIYAMA T. & NISHIHIRA J. Identification of macrophage migration inhibitory factor in murine neonatal calvariae and osteoblasts, 430
OROZCOE H. *see* HERNÁNDEZ-PANDO R.
OSIPOVA L. *see* ENGSTRÖM P.-E.

PAGE C.S. *see* MCDOUALL R.M.
PANAYOTOU G. *see* KLONISCH T.
PARKHOUSE R.M.E. *see* YANG H.
PARTIDOS C.D., PIZZA M., RAPPUOLI R. & STEWARD M.W. The adjuvant effect of a non-toxic mutant of heat-labile enterotoxin of *Escherichia coli* for the induction of measles virus-specific CTL responses after intranasal co-immunization with a synthetic peptide, 483
PASVOL G. *see* SANCHEZ-GARCIA J.
PAVÓN L. *see* HERNÁNDEZ-PANDO R.
PECORARO M.R., SHIMOJIMA M., MAEDA K., INOSHIMA Y., KAWAGUCHI Y., KAI C. & MIKAMI T. Molecular cloning of the feline CD8β chain, 84
PENNEY L., KILSHAW P.J. & MACDONALD T.T. Increased division of αβ TCR^+ and γδ TCR^+ intestinal intraepithelial lymphocytes after oral administration of cholera toxin, 54
PÉREZ C., CALDERÓN G.M., XIMÉNEZ C. & MELENDRO E.I. Human cell-mediated immune responses to antigenic fractions of *Salmonella typhi*, 262
PÉREZ-VILLAR J.J., ZAPATA J.M., MELERO I., POSTIGO A., SÁNCHEZ-MADRID F. & LÓPEZ-BOTET M. Expression and function of α_4/β_7 integrin on human natural killer cells, 96
PERRY F.E. *see* SHEN C.-R.
PERSSON M.A.A. *see* LUNDIN K.
PIETERS R. *see* ALBERS R.
PILLING D. *see* GOMBERT W.
PIÑEIRO A. *see* MARTÍNEZ-LORENZO M.J.
PIZZA M. *see* PARTIDOS C.D.
PORRECA E., REALE M., DI FEBBO C., DI GIOACCHINO M., BARBACANE R.C., CASTELLANI M.L., BACCANTE G., CONTI P. & CUCCURULLO F. Down-regulation of cyclooxygenase-2 (COX-2) by interleukin-1 receptor antagonist in human monocytes, 424
POSTIGO A. *see* PÉREZ-VILLAR J.J.
PROP J. *see* DE HAAN A.

QUENTMEIER H., DUSCHL A., HU Z.-B., SCHNARR B., ZABORSKI M. & DREXLER H.G. MUTZ-3, a monocytic model cell line for interleukin-4 and lipopolysaccharide studies, 606

RAMOS A., GONZÁLEZ M., LÓPEZ-HOYOS M., CARRIÓ R. & MERINO J. Completely allogeneic spleen cells induced cytolytic neonatal tolerance to alloantigens, but failed to establish allo-helper interactions with host T cells, 413
RAPPUOLI R. *see* PARTIDOS C.D.
RATH S. *see* MENDIRATTA S.K.
REALE M. *see* PORRECA E.
REEN D.J. *see* MURPHY F.J.
REES A.D.M. *see* FIDLER S.J.
REID H.W. *see* HAIG D. McK.
REILING N. *see* FLEISCHER J.
REINHOLD U., LIU L., SESTERHENN J. & ABKEN H. CD7-negative T cells represent a separate differentiation pathway in a subset of post-thymic helper T cells, 391
RESKE-KUNZ A.B. *see* FROSCH S.
RETZLAFF C., YAMAMOTO Y., OKUBO S., HOFFMAN P.S., FRIEDMAN H. & KLEIN T.W. *Legionella pneumophila* heat-shock protein-induced increase of interleukin-1β mRNA involves protein kinase C signalling in macrophages, 281
RINK L. *see* ALTSTAEDT J.
ROβ R. *see* FROSCH S.
ROBBINS R.A. *see* BERKMAN N.
ROBICHAUD A. *see* BERKMAN N.
ROESEMS G. *see* BERKMAN N.
ROITT I.M. *see* KLONISCH T.
ROLSTAD B. *see* HELGELAND L.
ROSATI E., SCARINGI L., CORNACCHIONE P., FETTUCCIARI K., SABATINI R., MEZZASOMA L., BENEDETTI C., CIANETTI S., ROSSI R. & MARCONI P. Activation of cytokine genes during primary and anamnestic immune response to inactivated *C. albicans*, 142
ROSE M.L. *see* MCDOUALL R.M.
ROSSI R. *see* ROSATI E.
ROWLAND S. *see* COATES J.P.

SABATINI R. *see* ROSATI E.
SAGAWA T. *see* OKADA M.
SAHEKI S. *see* HITSUMOTO Y.
SAITO H. *see* YANAGIDA M.
SALMON M. *see* GOMBERT W.
SAMPIERI A. *see* HERNÁNDEZ-PANDO R.
SAMUELSSON A. *see* LUNDIN K.
SANCEAU J. *see* LE PAGE C.
SANCHEZ-GARCIA J., ATKINS C., PASVOL G., WILKINSON R.J. & COLSTON M.J. Antigen-driven shedding of L-selectin from human $\gamma\delta$ T cells, 213
SÁNCHEZ-MADRID F. *see* PÉREZ-VILLAR J.J.
SAUDER D.N. *see* FUJISAWA H.
SAUL A.J. *see* BRANDT E.R.
SCARINGI L. *see* ROSATI E.
SCHLOTTMANN K. *see* GULBINS E.
SCHNARR B. *see* QUENTMEIER H.
SCHOENFELD D. *see* ISRAEL E.J.
SCHWING J. *see* FROSCH S.
SEINEN W. *see* ALBERS R.
SEKI S. *see* KAWAMURA T.
SELMAYR M. *see* KARDINAL C.
SEOW H.-F. *see* EGAN P.J.
see also HAIG D. McK.
SESTERHENN J. *see* REINHOLD U.
SHEN C.-R., MAZZA G., PERRY F.E., BEECH J.T., THOMPSON S.J., CORATO A., NEWTON S., BARKER R.N. & ELSON C.J. T-helper 1 dominated responses to erythrocyte Band 3 in NZB mice, 195
SHIMADA S. *see* KAWAGUCHI-MIYASHITA M.
SHIMIZU K. *see* KAWAGUCHI-MIYASHITA M.
SHIMOJIMA M. *see* PECORARO M.R.
SHINOMIYA T. *see* SUZUKI S.
SHIVJI G.M. *see* FUJISAWA H.
SILVA M.T. *see* LEAL I.S.
SIMISTER N.E. *see* ISRAEL E.J.
SINGH N. *see* MENDIRATTA S.K.
SOETH E. *see* FLEISCHER J.
SONG D.-S. *see* HITSUMOTO Y.
SORG R.V. *see* WILLIAMS L.A.
STEWARD M.W. *see* PARTIDOS C.D.
SUMMERS K.L. *see* WILLIAMS L.A.
SUZUKI K. *see* ONODERA S.
SUZUKI S., LI X.-K., ENOSAWA S. & SHINOMIYA T. A new immunosuppressant, FTY720, induces *bcl-2*-associated apoptotic cell death in human lymphocytes, 518
SYMON F.A. *see* WALSH G.M.

TADOKORO C.E., MACEDO M.S. & ABRAHAMSOHN I.A. Saponin adjuvant primes for a dominant interleukin-10 production to ovalbumin and to *Trypanosoma cruzi* antigen, 368
TAGUCHI O. & TAKAHASHI T. Mouse models of autoimmune disease suggest that self-tolerance is maintained by unresponsive autoreactive T cells, 13
TAKADA G. *see* KAWAMURA T.
TAKAHASHI T. *see* TAGUCHI O.
TAKEI M. *see* YANAGIDA M.
TAKENOYAMA M. *see* TAMADA K.
TAKEUCHI N. *see* HITSUMOTO Y.
TAMADA K., HARADA M., ITO O., TAKENOYAMA M., MORI T., MATSUZAKI G. & NOMOTO K. The emergence of non-cytolytic NK1.1$^+$ cells in the long-term culture of murine tumour-infiltrating lymphocytes: a possible role of transforming growth factor-β, 627
TAZAWA Y. *see* KAWAMURA T.
THIELEMANS K. *see* MURAILLE E.
THOMPSON S.J. *see* BARKER R.N.
see also SHEN C.-R.
TOKIWA T. *see* YANAGIDA M.
TOMINAGA A. *see* OKADA M.
TSUCHIYA S. *see* MURAI M.
TSUKADA T. *see* MIGITA K.
TSUKAMOTO H. *see* UEDA Y.

UEDA Y., NAGASAWA K., TSUKAMOTO H., HORIUCHI T., NISHIZAKA H., IKEDA K. & NIHO Y. Production of the third and fourth component of complement (C3, C4) by smooth muscle cells, 183
ULMER J.B., DECK R.R., DEWITT C.M., DONNELLY J.J. & LIU M.A. Generation of MHC class I-restricted cytotoxic T lymphocytes by expression of a viral protein in muscle cells: antigen presentation by non-muscle cells, 59
URBAIN J. *see* MURAILLE E.
USAMI J. *see* HIROMATSU K.
UZUMAKI H. *see* YANAGIDA M.

VAAGE J.T. *see* HELGELAND L.
VAINIO O. *see* KATEVUO K.
VAN DE WINKEL, J.G.J. *see* KRUGER M.
VAN DER PIJL A. *see* ALBERS R.
VAN MEURS M. *see* DEMAN J.
VAN ROOIJEN N. *see* DE HAAN A.
VELASQUILLO C. *see* HERNÁNDEZ-PANDO R.
VILLIERS C.L. *see* VILLIERS M.-B.
VILLIERS M.-B., VILLIERS C.L., JACQUIER-SARLIN M.R., GABERT F.M., JOURNET A.M. & COLOMB M.G. Covalent binding of C3b to tetanus toxin: influence on uptake/internalization of antigen by antigen-specific and non-specific B cells, 348

WAHREN B. *see* LUNDIN K.
WALLACE D.L. *see* GOMBERT W.
WALSH G.M., SYMON F.A., LAZAROVITS A.I. & WARDLAW A.J. Intergin α_4/β_7 mediates human eosinophil interaction with MAdCAM-1, VCAM-1 and fibronectin, 112
WANG B. *see* FUJISAWA H.
WANG Y. & YEH M. Molecular characterization of the V regions of four IgE antibodies specific for trichosanthin, 316
WARDLAW A.J. *see* WALSH G.M.
WATANABE T. *see* KAWAGUCHI-MIYASHITA M.
WEBER J. *see* FIDLER S.J.
WECHSLER-REYA R.J. & MONROE J.G. Lipopolysaccharide prevents apoptosis and induces responsiveness to antigen receptor cross-linking in immature B cells, 356

WEERASINGHE A. *see* KAWAMURA T.
WELLS A.D. *see* BARKER R.N.
WELSCH J. *see* GULBINS E.
WIEBE V. *see* ENGSTRÖM P.-E.
WIETZERBIN J. *see* LE PAGE C.
WIGZELL H. *see* LUNDIN K.
WILKINSON R.J. *see* SANCHEZ-GARCIA J.
WILLIAMS L.A., MCLELLAN A.D., SUMMERS K.L., SORG R.V., FEARNLEY D.B. & HART D.N.J. Identification of a novel dendritic cell surface antigen defined by carbohydrate specific CD24 antibody cross-reactivity, 120
WILLIAMS N.A. *see* HARPER H.M.
WILSCHUT J. *see* DE HAAN A.
WILSKER D.F. *see* ISRAEL E.J.
WOOD Y. *see* BRANDT E.R.

XIMÉNEZ C. *see* PÉREZ C.

YACOUB M.H. *see* MCDOUALL R.M.
YAGITA H. *see* FROSCH S.
YAMAMOTO Y. *see* RETZLAFF C.
YANAGIDA M., FUKAMACHI H., TAKEI M., HAGIWARA T., UZUMAKI H., TOKIWA T., SAITO H., IIKURA Y. & NAKAHATA T. Interferon-γ promotes the survival and FcεRI-mediated histamine release in cultured human mast cells, 547
YANG H. & PARKHOUSE R.M.E. Phenotypic classification of porcine lymphocyte subpopulations in blood and lymphoid tissues, 76
YEH M. *see* WANG Y.
YOSHIKAI Y. *see* HIROMATSU K.

ZABORSKI M. *see* QUENTMEIER H.
ZAPATA J.M. *see* PÉREZ-VILLAR J.J.

Subject Index

Allostimulation of human T cells by heart endothelial cells, 220
Anti-idiotic antibody response, autologous:
  regulation by idiotype, 172
Antigen binding:
  enhancement by synergy and antibody capture, 165
Antigen expression in muscle:
  generation of cytotoxic T lymphocyte, 59
Antigen-presenting cell defect, in HIV-1-infected patients:
  correlation with CD4 dependency in human T-cell clones, 46
Apoptosis:
  prevention by fibroblasts, 397
Apoptosis, *bcl-2*-associated:
  induced by FTY720, 518
Apoptosis, CD3 mediated:
  enhancement by antigen cross-linkage, 200
Autoantigens:
  in pristane-induced arthritis, 189
Autoimmunity:
  chemical induction of, 468

B cells, human:
  in vitro immunization, 1

C3b:
  covalent binding to tetanus toxin, 348
CD7-negative T cells, 391
CD14 expression:
  effect of granulocyte–macrophage colony-stimulating factor, 89
CD24 monoclonal antibody:
  cross-reactivity on dendritic cells, 120
CD8β, feline:
  molecular cloning, 84
CD80 and CD86:
  differential expression and function on human monocytes, 592
CD88:
  expression and modulation on skin dendritic cells, 126
CDR3 sequences, IgH chain:
  human B-lineage cells, 324
Cell-mediated immunity to *Salmonella typhi* antigens, 262
Complement C3 and C4 production:
  by smooth muscle cells, 183
Complement regulatory protein:
  expression by human oligodendrocyte cell line, 338
Costimulatory signalling:
  by major histocompatibility complex class II^+ T cells, 384
Cyclo-oxygenase-2:
  down-regulation by interleukin-2 receptor antagonist, 424
Cyclosporin A:
  abrogation of acquired immunity to parapoxvirus orf virus, 524
Cytokine genes:
  activation in responses to *Candida albicans*, 142
Cytokine profiles:
  in CD4 knockout mice, 250
Cytotoxic T-cell responses:
  requirement for $CD4^+$ T-cell help, 41
Cytotoxicity:
  of intermediate T-cell receptor, 68

Dendritic cells:
  phenotype and function, 457
Diabetes, in NOD mice:
  prevention by Mycobacterial infection, 20

Fas ligand, release of preformed:
  by Jurkat T cells, 511

Ficolin, human:
  biosynthesis in monocytes, 289

IgG clearance:
  in β_2-microglobulin deficient mice, 573
Immunoglobulin loci:
  gene targetting, 309
Immunoglobulins, duck:
  ontogeny of, 8
Inducible nitric oxide synthase:
  in lung epithelial cells, 363
Inflammation:
  induced changes in migrating cells, 539
Integrin α_4/β_7, on natural killer cells:
  function, 96
Integrin α_4/β_7:
  mediation of eosinophil interactions, 112
β7 integrins:
  in human gut, 227
Interleukin-1β mRNA, induced by heat-shock protein:
  role of protein kinase C signalling, 281
Interleukin-2 response, to superantigens:
  regulation by costimulation, 245
Interleukin-6 production:
  after infection with *Schistosoma mansoni*, 256
Interleukin-6 receptor:
  functional epitope, 135
Interleukin-7:
  influence on V(D)J recombination, 569
Interleukin-10 production:
  priming by saponin adjuvant, 368
Intraepithelial lymphocytes, intestinal:
  effect of cholera toxin, 54
Intraepithelial lymphocytes:
  in antigen-minimized mice, 268
  T-cell receptor Vβ repertoire, 494
Intranasal immunization:
  induction of CTL with *Escherichia coli* heat-labile enterotoxin as adjuvant, 483

L-selectin:
  from human $\gamma\delta$ T cells, 213
Lipids, from *Mycobacterium leprae* cell wall:
  anti-inflammatory property, 613
Lipopolysaccharide:
  and responsiveness in immature B cells, 356

α_2-macroglobulin:
  stimulation of protein tyrosine kinase, 436
Macrophage migration inhibitory factor:
  in murine calvariae and osteoblasts, 430
Macrophages, alveolar:
  role in liposome adjuvant activity, 488
Major histocompatibility complex class II expression, on monocytes:
  effect of lipopolysaccharide, 599
Mast cells, human:
  FcεRI-mediated histamine release, 547
Mast cells, serotonin release:
  antigen triggered, 463
Monocyte subpopulation:
  function-related antigens, 587
Monocytic model cell line:
  for interleukin-4 and lipopolysaccharide studies, 606

Neutrophil priming, 105

Neutrophils:
in murine listeriosis, 442
production of interleukin-8, 563
Nitric oxide:
role in activity of rat neutrophils, 295
Nitric oxide synthase:
differential expression, 274

Opsonic antibodies:
specific for M protein of group A streptococci, 331
Oral tolerance:
intestinal lamina propria, 449

Peptide library:
immunological mimicry of a neutralizing HIV-1 epitope, 579
Platelet-mediated tumour cell killing:
via cyclo-oxygenase or nitric oxide pathway, 158
Porcine lymphocytes:
classification, 76
Programmed cell death, Fas induced:
mediated by Ras-regulated O_2^- synthesis, 205
Prolactin:
effect on interleukin-2-driven lymphokine-activated killed cells, 619

Salivary IgG subclasses:
and IGHG deletions, 178
Self-tolerance:
maintenance by unresponsive T cells, 13
Stem cell factor:
modulation of mast cell histamine and tumour necrosis factor-α secretion, 301

T-cell hybridomas:
major histocompatibility complex class II ligand specificity, 238
T-cell subsets:
in anterior chamber immune deviation, 34
in chemical sensitization, 502
γδ T cells:
response to *Plasmodium falciparum*, 405
T-helper 1 cytokine responses:
in NZB mice, 195
T-helper cell subsets, kinetics:
and pathology in experimental tuberculosis, 26
Th0-CD4 T-cell responses:
in murine retrovirus-induced immunodeficiency syndrome, 532
Thymocyte emigration:
chicken, 419
Thymus, fetal:
in vivo cytokine expression, 152
Tolerance, neonatal:
allogeneic B cells, 413
Tumour necrosis factor-α production:
inhibition by 3-deazaadenosine, 558
Tumour-infiltrating lymphocytes:
emergence of non-cytolytic $NK1.1^+$ T cells, 627
Tumour-necrosis factor-α-mediated expression of membrane-type matrix metalloproteinase, 553

V region genes:
of IgE antibodies, 316
Vascular cell adhesion molecule-1 expression:
by fibroblasts, 375